

A *rational function* is a function of the form

$$f(x) = \frac{g(x)}{h(x)},$$

where $g(x)$ and $h(x)$ are polynomials. A rational function is in *lowest form* if the numerator and the denominator have no common zeros. Assume that $f(x) = g(x)/h(x)$ is a rational function in lowest form.

The *degree* of $f(x)$ is $\max\{\deg(g), \deg(h)\}$.

The *roots* of $f(x)$ are the zeros of $g(x)$; that is, they are the solutions to $g(x) = 0$.


The *poles* of $f(x)$ are the zeros of $h(x)$; that is, they are the solutions to $h(x) = 0$.

The *y-intercept* of $f(x)$ is the point $(0, f(0))$.

The *x-intercepts* of $f(x)$ are the points $(r, 0)$, where r is a root of $f(x)$.

The *vertical asymptotes* of $f(x)$ are the lines $x = p$, where p is a pole of $f(x)$.

The *polynomial asymptote* of $f(x)$ is the polynomial equation $y = q(x)$, where $q(x)$ is the quotient when $g(x)$ is divided by $h(x)$ using polynomial division.



Problem 1: $f(x) = \frac{2}{x-1}$

Degree:

Roots:


Poles:

y-intercept:

x-intercepts:

Vertical Asymptotes:

Polynomial Asymptote:



Problem 2: $f(x) = \frac{6x+3}{2x-4}$

Degree:

Roots:


Poles:

y-intercept:

x-intercepts:

Vertical Asymptotes:

Polynomial Asymptote:



Problem 3: $f(x) = \frac{x^2 - 2x - 15}{x + 1}$

Degree:

Roots:


Poles:

***y*-intercept:**

***x*-intercepts:**

Vertical Asymptotes:

Polynomial Asymptote:



Problem 4: $f(x) = \frac{x^2 - 49}{x^2 - 25}$

Degree:

Roots:


Poles:

***y*-intercept:**

***x*-intercepts:**

Vertical Asymptotes:

Polynomial Asymptote:



Problem 5: $f(x) = \frac{x^3 - x}{x^2 - 9}$

Degree:

Roots:

Poles:

***y*-intercept:**

***x*-intercepts:**

Vertical Asymptotes:

Polynomial Asymptote: