

## THE COSINE OF $72^\circ$

PAUL L. BAILEY

**Problem 1.** Compute the cosine of  $72^\circ$ .

*Solution via Trigonometry.* Let  $\theta = 18^\circ$ ; then  $\sin \theta = \cos 72^\circ$ . Now  $5\theta = 90^\circ$ , so  $3\theta = 90^\circ - 2\theta$ ; therefore

$$\sin 3\theta = \cos 2\theta.$$

Use the sine sum of angles formula on the left:

$$\sin 2\theta \cos \theta + \sin \theta \cos 2\theta = \cos 2\theta.$$

Subtract  $\sin \theta \cos 2\theta$  from both sides:

$$\sin 2\theta \cos \theta = \cos 2\theta(1 - \sin \theta).$$

Use the sine double angle formula:

$$2 \sin \theta \cos^2 \theta = \cos 2\theta(1 - \sin \theta).$$

Apply the cosine squared and cosine double angle identities:

$$2 \sin \theta(1 - \sin^2 \theta) = (1 - 2 \sin^2 \theta)(1 - \sin \theta).$$

Divide by  $(1 - \sin \theta)$  to get

$$2 \sin \theta(1 + \sin \theta) = 1 - 2 \sin^2 \theta.$$

Make a polynomial in  $\sin \theta$ :

$$4 \sin^2 \theta + 2 \sin \theta - 1 = 0.$$

Apply the quadratic formula to find that

$$\begin{aligned} \sin \theta &= \frac{-2 \pm \sqrt{4 + 16}}{8} \\ &= \frac{-1 \pm \sqrt{5}}{4}. \end{aligned}$$

Since  $18^\circ$  is in the first quadrant, the sine must be positive; we conclude that

$$\cos 72^\circ = \sin 18^\circ = \frac{-1 + \sqrt{5}}{4}.$$

□

*Solution via Complex Numbers.* First we note that  $72^\circ = \frac{360^\circ}{5} = \frac{\pi}{5}$  radians.

We wish to use complex numbers on the unit circle to solve this problem, Let  $\alpha = e^{2\pi i/5}$ ; we wish to find  $\cos \frac{\pi}{5} = \Re \alpha$ .

The complex number  $\alpha$  is a primitive 5<sup>th</sup> root of unity, and so  $\alpha$  satisfies the polynomial equation  $x^5 - 1 = 0$ . Clearly 1 is a root of  $x^5 - 1$ ; factor this out to obtain  $x^5 - 1 = (x - 1)(x^4 + x^3 + x^2 + x + 1)$ . Now  $\alpha$  is a root of the latter factor, so  $\alpha^4 + \alpha^3 + \alpha^2 + \alpha + 1 = 0$ .

Note that  $\alpha^4$  is the complex conjugate of  $\alpha$  and  $\alpha^2$  is the complex conjugate of  $\alpha^3$ . Thus,  $\Re \alpha = \frac{1}{2}(\alpha + \alpha^4)$  and  $\Re \alpha^2 = \frac{1}{2}(\alpha^2 + \alpha^3)$ . Let  $\zeta_1 = (\alpha + \alpha^4)$  and  $\zeta_2 = (\alpha^2 + \alpha^3)$ ; we have  $\zeta_1 + \zeta_2 = -1$ . Compute

$$\zeta_1 \zeta_2 = \alpha^3 + \alpha^4 + \alpha^6 + \alpha^7 = \alpha^3 + \alpha^4 + \alpha + \alpha^2 = -1.$$

Define the quadratic function

$$f(x) = (x - \zeta_1)(x - \zeta_2) = x^2 - (\zeta_1 + \zeta_2)x + \zeta_1 \zeta_2 = x^2 + x - 1.$$

The roots of  $f$  are given by the quadratic formula to be

$$x = \frac{-1 \pm \sqrt{5}}{2}.$$

Geometrically, it is clear that  $\zeta_1 > 0$  and  $\zeta_2 < 0$ ; thus

$$\zeta_1 = \frac{-1 + \sqrt{5}}{2} \quad \text{and} \quad \zeta_2 = \frac{-1 - \sqrt{5}}{2}.$$

Thus

$$\cos 72^\circ = \Re \alpha = \frac{-1 + \sqrt{5}}{4}.$$

□

*Solution via the Golden Triangle.* Consider an isosceles triangle  $\triangle ABC$  with the property that the equal angles  $\angle ABC$  and  $\angle ACB$  are twice the other angle  $\angle BAC$ . Let  $\alpha = \angle BAC$  and  $\beta = \angle ABC = \angle ACB$ , so that  $\beta = 2\alpha$ . Then  $5\alpha = 180^\circ$ , so  $\alpha = 36^\circ$  and  $\beta = 72^\circ$ . We wish to find  $\cos \beta$ .

Bisect  $\angle ABC$  and let  $D$  be the point on  $\overline{AC}$  such that  $\angle BCD = \alpha$ . Now  $\triangle DAB$  is isosceles, with  $|DA| = |BA|$ , and  $\triangle BCD$  is similar to  $\triangle ABC$ . Let  $x = |AB|$ ,  $y = |BC|$ , and  $z = |CD|$ , so that  $x = y + z$ . Moreover,  $\frac{x}{y} = yz$  by similarity. Thus  $y^2 = xz = (y + z)z = yz + z^2$ , so  $y^2 - yz - z^2 = 0$ , and by the quadratic formula,

$$y = \frac{z + \sqrt{z^2 + 4z^2}}{2} = z \frac{1 + \sqrt{5}}{2}.$$

Thus

$$\cos \beta = \frac{z}{2y} = \frac{1}{1 + \sqrt{5}} = \frac{-1 + \sqrt{5}}{5 - 1}.$$

Conclude that

$$\cos 72^\circ = \frac{-1 + \sqrt{5}}{4}.$$

□