Vector Calculus Quadric Surface Classification Dr. Paul L. Bailey Wednesday, September 18, 2019

A quadric surface is the locus of an equation of the form

 $Ax^{2} + By^{2} + Cz^{2} + Dx + Ey + Fz + Gxy + Hxz + Iyz = J.$

We may eliminate the mixed terms by rotations; thus assume that the equation is of the form

 $Ax^2 + By^2 + Cz^2 + Dx + Ey + Fz = J.$

In the presence of a quadratic (degree 2) term we may eliminate the linear (degree 1) term via completing the square to obtain a translation; thus assume that each variable occurs as a quadratic or as a linear term, but not both.

We call J the *constant*, and assume that $J \ge 0$.

A *slice* is the intersection of a quadric surface with a plane perpendicular to a coordinate plane. We classify the surfaces as follows.

- If all variables are linear, it is a *plane*.
- If one variable is missing, it is a *cylinder*.
- If two variables are linear, it is a *cylinder*.
- If two variables are quadratic and one is linear, it is a *paraboloid*. In this case, two slices are parabolas.
 - If the quadratics have the same sign, it is an *elliptic paraboloid*.
 - If the quadratics have different signs, it is a hyperbolic paraboloid, or "saddle".
- If three variables are quadratic and positive, it is an *ellipsoid*.
- If three variable are quadratic and one has a different sign, and the constant is zero, it is a *cone*.
- If three variables are quadratic and one has a different sign, and the constant is positive, it is a *hyperboloid*. In this case, two slices are hyperbolas.
 - If two quadratics are positive, it is a one-sheeted hyperboloid.
 - If two quadratics are negative, it is a two-sheeted hyperboloid.

We give an example equation for each type.

• Plane	x + y + z = 1
• Cylinder	$z = x^2 + y$
• Elliptic Paraboloid	$z = x^2 + y^2$
• Hyperbolic Paraboloid	$z = x^2 - y^2$
• Ellipsoid	$x^2 + y^2 + z^2 = 1$
• One-Sheeted Hyperboloid	$x^2 + y^2 - z^2 = 1$
• Two-Sheeted Hyperboloid	$z^2 - x^2 - y^2 = 1$
• Cone	$x^2 + y^2 - z^2 = 0$