

# Trigonometry Summary

## Dr. Paul L. Bailey

### Definitions

The *radian measure* of an angle is the arclength of the arc on a circle of radius 1 which subtends the angle when the vertex is placed at the center of the circle. We can convert between degrees and radians by using the fact that

$$360^\circ = 2\pi \text{ radians.}$$

The *wrapping function*  $P : \mathbb{R} \rightarrow \mathbb{R}^2$  is defined to be the point  $P(\theta)$  on the unit circle  $x^2 + y^2 = 1$  obtained by moving counterclockwise along the circle an arclength of  $\theta$ . If  $\theta < 0$ , this is interpreted as moving clockwise by an arclength of  $|\theta|$ .

The *sine* and *cosine* functions as defined by

$$\sin \theta = \text{the } y\text{-coordinate of } P(\theta) \quad \text{and} \quad \cos \theta = \text{the } x\text{-coordinate of } P(\theta).$$

It is clear that

$$\sin^2 \theta + \cos^2 \theta = 1.$$

The *tangent*, *cotangent*, *secant*, and *cosecant* are defined by

$$\tan \theta = \frac{\sin \theta}{\cos \theta}, \quad \cot \theta = \frac{\cos \theta}{\sin \theta}, \quad \sec \theta = \frac{1}{\cos \theta}, \quad \csc \theta = \frac{1}{\sin \theta}.$$

### Trigonometric Identities

#### Identities that Come from Geometry

Description	Identity
<b>Pythagorean Identity</b>	$\cos^2 \theta + \sin^2 \theta = 1$
<b>Symmetry</b>	$\sin(-\theta) = -\sin \theta$ $\cos(-\theta) = \cos \theta$
<b>Periodicity</b>	$\sin(\theta + 2\pi) = \sin \theta$ $\cos(\theta + 2\pi) = \cos \theta$
<b>Rotation by 180°</b>	$\sin(\theta + \pi) = -\sin \theta$ $\cos(\theta + \pi) = -\cos \theta$
<b>Rotation by 90°</b>	$\sin(\theta + \frac{\pi}{2}) = \cos \theta$ $\cos(\theta - \frac{\pi}{2}) = \sin \theta$
<b>Reflection across 45°</b>	$\sin(\frac{\pi}{2} - \theta) = \cos \theta$ $\cos(\frac{\pi}{2} - \theta) = \sin \theta$
<b>Difference of Angles Formula</b>	$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$

#### Identities that Come from Algebra

Description	Identity
<b>Sum and Difference Formulas</b>	$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$ $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$ $\sin(\alpha + \beta) = \sin \alpha \cos \beta + \sin \beta \cos \alpha$ $\sin(\alpha - \beta) = \sin \alpha \cos \beta - \sin \beta \cos \alpha$
<b>Double Angle Formulas</b>	$\cos(2\theta) = \cos^2 \theta - \sin^2 \theta$ $\sin(2\theta) = 2 \sin \theta \cos \theta$
<b>Half Angle Formulas</b>	$\sin \frac{\theta}{2} = \sqrt{\frac{1 - \cos \theta}{2}}$ $\cos \frac{\theta}{2} = \sqrt{\frac{1 + \cos \theta}{2}}$
<b>Pythagorean Formulae</b>	$1 + \tan^2 \theta = \sec^2 \theta$ $\cot^2 \theta + 1 = \csc^2 \theta$

### Trigonometric Values

deg( $\theta$ )	rad( $\theta$ )	sin( $\theta$ )	cos( $\theta$ )	tan( $\theta$ )	cot( $\theta$ )	sec( $\theta$ )	csc( $\theta$ )
0°	0	0	1	0	$\infty$	1	$\infty$
15°	$\frac{\pi}{12}$	$\frac{\sqrt{6} - \sqrt{2}}{4}$	$\frac{\sqrt{6} + \sqrt{2}}{4}$	$\frac{2 - \sqrt{3}}{2}$	$\frac{2 + \sqrt{3}}{2}$	$\sqrt{6} - \sqrt{2}$	$\sqrt{6} + \sqrt{2}$
18°	$\frac{\pi}{10}$	$\frac{\sqrt{5} - 1}{4}$	$\frac{\sqrt{10 + 2\sqrt{5}}}{4}$	$\frac{5 - 2\sqrt{5}}{2}$	$\sqrt{5 + 2\sqrt{5}}$	$\sqrt{2\sqrt{5} - 5}$	$\sqrt{5} + 1$
	$\frac{\pi}{8}$	$\frac{\sqrt{2 - \sqrt{2}}}{2}$	$\frac{\sqrt{2 + \sqrt{2}}}{2}$	$\sqrt{3 - 2\sqrt{2}}$	$\sqrt{3 + 2\sqrt{2}}$	$\sqrt{4 - 2\sqrt{2}}$	$\sqrt{4 + 2\sqrt{2}}$
30°	$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$	$\sqrt{3}$	$\frac{2\sqrt{3}}{3}$	2
36°	$\frac{\pi}{5}$	$\frac{\sqrt{10 - 2\sqrt{5}}}{4}$	$\frac{1 + \sqrt{5}}{4}$	$\sqrt{5 - 2\sqrt{5}}$	$\frac{\sqrt{25 + 10\sqrt{5}}}{5}$	$\sqrt{5} - 1$	$\frac{10 + 2\sqrt{5}}{5}$
45°	$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	1	$\sqrt{2}$	$\sqrt{2}$
54°	$\frac{3\pi}{10}$	$\frac{1 + \sqrt{5}}{4}$	$\frac{\sqrt{10 - 2\sqrt{5}}}{4}$	$\frac{\sqrt{25 + 10\sqrt{5}}}{5}$	$\sqrt{5 - 2\sqrt{5}}$	$\frac{10 + 2\sqrt{5}}{5}$	$\sqrt{5} - 1$
60°	$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2\sqrt{3}}{3}$
	$\frac{3\pi}{8}$	$\frac{\sqrt{2 + \sqrt{2}}}{2}$	$\frac{\sqrt{2 - \sqrt{2}}}{2}$	$\sqrt{3 + 2\sqrt{2}}$	$\sqrt{3 - 2\sqrt{2}}$	$\sqrt{4 + 2\sqrt{2}}$	$\sqrt{4 - 2\sqrt{2}}$
72°	$\frac{2\pi}{5}$	$\frac{\sqrt{10 + 2\sqrt{5}}}{4}$	$\frac{\sqrt{5} - 1}{4}$	$\sqrt{5 + 2\sqrt{5}}$	$\frac{5 - 2\sqrt{5}}{2}$	$\sqrt{5} + 1$	$\sqrt{2\sqrt{5} - 5}$
75°	$\frac{5\pi}{12}$	$\frac{\sqrt{6} + \sqrt{2}}{4}$	$\frac{\sqrt{6} - \sqrt{2}}{4}$	$\frac{2 + \sqrt{3}}{2}$	$\frac{2 - \sqrt{3}}{2}$	$\sqrt{6} + \sqrt{2}$	$\sqrt{6} - \sqrt{2}$
90°	$\frac{\pi}{2}$	1	0	$\infty$	0	$\infty$	1